

Universal Properties of Light Rings for Stationary Axisymmetric Spacetimes

Speaker : Sijie Gao 高思杰
(北京师范大学 Beijing Normal University)

Cooperated with Minyong Guo 郭敏勇
(北京大学 Peking University)

arXiv:2011.02211 (To be published in PRD)

2021年5月6日 Tsinghua University

1.Motivation

- A light ring (LR) is a circular photon orbit outside a black hole or an ultracompact object (UCO)
- Light ring is an important feature of curved spacetimes.
- Light rings play an important role in gravitational wave observations and black hole photographs.
- Light rings could be observational evidence for event horizons [V. Cardoso, et.al. PRL(2014)]

2.Example: Schwarzschild black hole

The metric for the Schwarzschild black hole is:

$$ds^2 = \left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2$$

Due to the spherical symmetry, we can always assume that the light moves on the equatorial plane $\theta = \pi/2$. The four-momentum is

$$p^a = \dot{t} \left(\frac{\partial}{\partial t}\right)^a + \dot{r} \left(\frac{\partial}{\partial r}\right)^a + \dot{\phi} \left(\frac{\partial}{\partial \phi}\right)^a$$

where $\dot{t} = \frac{\partial t}{\partial \lambda}$ and λ is an affine parameter of the null geodesic.

There are two conserved quantities, energy and angular momentum associated with the two Killing vector fields $\left(\frac{\partial}{\partial t}\right)^a$ and $\left(\frac{\partial}{\partial \phi}\right)^a$:

$$E = -g_{ab}p^a \left(\frac{\partial}{\partial t}\right)^b = (1 - 2M/r)\dot{t}$$

$$L = g_{ab}p^a \left(\frac{\partial}{\partial \phi}\right)^b = r^2\dot{\phi}$$

For null curves, we have

$$0 = g_{ab}p^a p^b = -(1 - 2M/r)\dot{t}^2 + (1 - 2M/r)^{-1}\dot{r}^2 + r^2\dot{\phi}^2$$

Thus, the radial geodesic equation can be solved as

$$\dot{r}^2 = E^2 - V(r)$$


with

$$V(r) = \frac{L^2}{r^2} - \frac{2ML^2}{r^3}$$


The light ring $r = r_0$ satisfies


$$\dot{r}^2 = E^2 - V(r_0) = 0$$

$$\left. \frac{\partial V}{\partial r} \right|_{r=r_0} = 0$$



The unstable light ring is located at $r = 3M$.







From Wald et al. 2019

Do the light rings exist in general
in curved spacetimes?

3. Previous works


- P. Cunha et.al shows that light rings always come in pairs for UCO (PRL 2017), one being a saddle point and the other a local extremum. A topological argument was used in the proof.



- Very recently, the authors employed this topological argument to a stationary black hole and found that at least one standard LR exists out-side the nonextremal horizon for each rotation sense (PRL 2020).

Unresolved issues

- The topological argument relies on the formation history of UCO
- The previous works cannot answer that whether a light ring must exist on the equatorial plane.



- It cannot answer whether the radial or the angular direction is stable.
- The topological argument does not apply to extremal black holes.

4. General setup

- A stationary spacetime with two Killing vector fields:

$$ds^2 = g_{tt}(r, \theta)dt^2 + g_{rr}(r, \theta)dr^2 + 2g_{t\phi}(r, \theta)dtd\phi + g_{\theta\theta}(r, \theta)d\theta^2 + g_{\phi\phi}(r, \theta)d\phi^2$$

- In general, the 4-momentum of a photon is

$$p^a = \dot{t} \left(\frac{\partial}{\partial t} \right)^a + \dot{r} \left(\frac{\partial}{\partial r} \right)^a + \dot{\theta} \left(\frac{\partial}{\partial \theta} \right)^a + \dot{\phi} \left(\frac{\partial}{\partial \phi} \right)^a$$

With $p^a p_a = 0$

- The conservation of energy and angular momentum:

$$E = -g_{ab}p^a \left(\frac{\partial}{\partial t} \right)^b = -g_{tt}\dot{t} - g_{t\phi}\dot{\phi},$$

$$L = g_{ab}p^a \left(\frac{\partial}{\partial \phi} \right)^b = g_{\phi\phi}\dot{\phi} + g_{t\phi}\dot{t},$$

• Therefore $g_{rr}\dot{r}^2 + g_{\theta\theta}\dot{\theta}^2 + V(r, \theta) = 0$

Where $V(r, \theta) = -\frac{1}{D} (E^2 g_{\phi\phi} + 2ELg_{t\phi} + L^2 g_{tt})$

$$D \equiv g_{t\phi}^2 - g_{tt}g_{\phi\phi}.$$

Define $\sigma = \frac{E}{L}$. Then

$$V = -\frac{L^2 g_{\phi\phi}}{D} (\sigma - H_+) (\sigma - H_-)$$

$$H_{\pm} = \frac{-g_{t\phi} \pm \sqrt{D}}{g_{\phi\phi}}.$$

For the H_+ branch, assume that a LR is located at $\{r = r_+, \theta = \theta_+\}$, which satisfies

$$\partial_r H_+ \Big|_{(r_+, \theta_+)} = 0, \quad \partial_\theta H_+ \Big|_{(r_+, \theta_+)} = 0, \quad \sigma = H_+(r_+, \theta_+)$$

Similarly for the H_- branch.

- The H_+ branch corresponds to LR counter-rotating with the black hole and the H_- branch corresponds to LR co-rotating with the black hole.
- To find LRs, we shall analyze the angular direction and the radial direction, respectively.

Stability

$$\partial_m^2 V(r_{\text{LR}}, \theta_{\text{LR}}) = \partial_m^2 H_+(r_{\text{LR}}, \theta_{\text{LR}})(H_+ - H_-) .$$

$$\partial_m^2 V(r_{\text{LR}}, \theta_{\text{LR}}) = \partial_m^2 H_-(r_{\text{LR}}, \theta_{\text{LR}})(H_- - H_+) .$$

where, $m \in \{r, \theta\}$

Since $H_+ > H_-$, we see that $\partial_m^2 H_+$ has the same sign as $\partial_m^2 V$ and $\partial_m^2 H_-$ has the opposite sign.

So the maximum of H_+ and the minimum of H_- correspond to unstable orbits.

5. Existence of LR for Black holes

- Axis-symmetric black holes--angular direction
(fixed r)


$\rho \equiv \sqrt{g_{\phi\phi}}$ goes to zero when $\theta \rightarrow 0$ and $\theta \rightarrow \pi$.

$$\begin{cases} \partial_{\theta}\rho > 0 & \theta \rightarrow 0, \\ \partial_{\theta}\rho < 0 & \theta \rightarrow \pi. \end{cases}$$

Thus we find $H_{\pm} \simeq \pm \frac{1}{\rho} \rightarrow \pm\infty,$

and $\partial_{\theta}H_{\pm} \sim \mp \frac{\partial_{\theta}\rho}{\rho^2} \sim \begin{cases} \mp\infty & \theta \rightarrow 0 \\ \pm\infty & \theta \rightarrow \pi, \end{cases}.$

So, for any fixed r , $H_{\pm}$ can be viewed as a function of θ ranging from 0 to π ,



For each given r , there always exists a $\theta = \theta_+$ such that $H_+(r, \theta_+)$ is a minimum in the θ direction. In this way, we obtain a function $\theta = \theta_+(r)$. Similarly, we have $\theta_-(r)$ for H_- .

In asymptotically flat spacetimes, $H_{\pm} \rightarrow \pm \frac{1}{r \sin \theta}$ as $r \rightarrow \infty$. Thus, we have

$$\theta_{\pm}(r \rightarrow \infty) = \pi/2.$$

When the spacetime possesses a parity reflection symmetry $\theta \rightarrow \pi - \theta$, H_+ satisfies $H_+(\theta) = H_+(\pi - \theta)$. Thus,

$$\left. \frac{\partial H_+}{\partial \theta} \right|_{\theta} = - \left. \frac{\partial H_+}{\partial \theta} \right|_{\pi - \theta}$$

On the equatorial plane,

$$\left. \frac{\partial H_+}{\partial \theta} \right|_{\pi/2} = - \left. \frac{\partial H_+}{\partial \theta} \right|_{\pi/2} = 0$$

Therefore, $\theta_{\pm}(r) = \pi/2$.

An orbit confined on the equatorial plane always has an extremum in the angular direction.

Radial direction (fixed θ)


- At infinity $r \rightarrow \infty$, we have

$$g_{t\phi} \rightarrow 0, \quad g_{tt} \rightarrow -1 \quad \text{and} \quad g_{\phi\phi} \rightarrow r^2$$

$$H_{\pm} \rightarrow \pm \frac{1}{r \sin \theta} \rightarrow 0^{\pm}$$

- On the horizon $D|_{r_h} = 0$, and D is always positive outside the horizon

$$H_{\pm}|_{r_h} = - \frac{g_{t\phi}}{g_{\phi\phi}} \Big|_{r_h} < 0.$$



So there exists at least one maximum for H_+

H_- is always negative outside the horizon.



$$H'_-(r_h) \sim -\frac{D'(r_h)}{2\sqrt{D(r_h)}g_{\phi\phi}(r_h)} + \frac{g_{t\phi}(r_h)g'_{\phi\phi}(r_h) - g'_{t\phi}(r_h)g_{\phi\phi}(r_h)}{g_{\phi\phi}^2(r_h)};$$

One can show that $D'(r_h) > 0$ for nonextremal horizons


Thus, $H'_-(r_h) \rightarrow -\infty$

There exists at least one minimum for H_- (maximum for V)


The above argument holds for any constant θ , i.e., given any θ , there exists $r = r_+ > r_h$ such that H_+ takes an maximum value in the r direction. Hence, we have a function $r_+(\theta)$ defined in the range $0 \leq \theta \leq \pi$.



The functions $r_{\pm}(\theta)$ and $\theta_{\pm}(r)$ for the Kerr black hole



- For extremal Kerr black holes $D'(r_h) = 0$



6. LRs in horizonless spacetime

- Horizonless spacetimes are interesting because they can represent ultracompact objects (UCO)
- The behavior of $H_{\pm}$ at infinity is the same as BH,

$$H_{\pm} \rightarrow 0^{\pm}$$

- Assume that the spacetime is regular,
 $g_{tt} < 0$ and $g_{t\phi} > 0$ everywhere.

- Near the center $r=0$, assume

$$g_{\phi\phi} \rightarrow r^2 \quad g_{tt} \rightarrow -k^2 \quad g_{t\phi} \rightarrow pr^s$$

$$H_{\pm} \sim \frac{-pr^s \pm \sqrt{p^2 r^{2s} + k^2 r^2}}{r^2}$$

For $s > 1$,
$$H_{\pm} \sim \frac{-pr^s \pm \sqrt{k^2 r^2}}{r^2} \sim \pm \frac{1}{r} \rightarrow \pm \infty.$$

For $s = 1$,
$$H_{\pm} \sim \frac{\pm \sqrt{p^2 + k^2} - p}{r} \rightarrow \pm \infty.$$

For $0 < s < 1$,

$$H_+ \sim \frac{-pr^s + pr^s (1 + \alpha^2 r^{2-2s})^{1/2}}{r^2} \sim \frac{\alpha^2 p}{2} r^{-s} \rightarrow \infty$$


$$H_- \sim - \left(\frac{2}{r^{2-s}} + \frac{\alpha^2}{2r^s} \right) p \rightarrow -\infty. \quad \alpha \equiv \frac{k}{p}.$$

for $s = 0$,
$$H_{\pm} \sim \frac{-p \pm \sqrt{p^2 + k^2 r^2}}{r^2} = \frac{-1 \pm \sqrt{1 + \alpha^2 r^2}}{r^2} p,$$

$$H_- \rightarrow -\infty \quad H_+ \rightarrow \frac{\alpha^2}{2} p > 0$$

$$H'_+(0^+) = 0, \quad \text{and} \quad H''_+(0^+) = -\frac{\alpha^4}{4} p < 0,$$

Hence, H'_+ become negative just away from $r = 0$



- There is either no LR or there are even number of LRs for each branch.

7. Conclusions

- We have shown that there are at least two LRs outside a non-extreme stationary black hole. The outermost ELRs are unstable along the radial direction and stable along the angular direction.
- For an extremal stationary black hole, we find there is at least one LR.
- When the spacetime possesses a reflection symmetry, there must exist a LR on the equatorial plane.

- For horizonless spacetimes, we have proved that if LR exists, there are at least two LRs, with the outer one being unstable in the radial direction and the inner one being stable.
- In our arguments, only some generic conditions have been used, for instance, the asymptotically flat condition and the behaviors of the metric near the horizon or the center of star. The results could apply to most gravity theories.

Thank You !